

# Groundstate properties of the unbound $T_z = 5/2$ nucleus $^{15}\text{Ne}^*$

F. Wamers<sup>1,2,3</sup>, J. Marganec<sup>2,1,3</sup>, T. Aumann<sup>2,1</sup>, L. Chulkov<sup>4,1</sup>, M. Heil<sup>1</sup>, B. Jonson<sup>5</sup>, T. Nilsson<sup>5</sup>,  
R. Plag<sup>1</sup>, H. Simon<sup>1</sup>, and the R<sup>3</sup>B Collaboration

<sup>1</sup>GSI, Darmstadt; <sup>2</sup>Technische Universität, Darmstadt; <sup>3</sup>EMMI @ GSI, Darmstadt; <sup>4</sup>NRC Kurchatov Institute, Moscow; <sup>5</sup>Chalmers Tekniska Högskola, Göteborg

In this report we present our findings on the properties of the recently observed  $^{15}\text{Ne}$  [1], with a focus on its ground-state structure and the decay mechanism to  $^{13}\text{O}$ .




Fig. 1: Level scheme of  $^{15}\text{Ne}$  and neighbours along its decay path to  $^{13}\text{O}$  [2]. Decay via  $^{14}\text{F}$  is energetically possible.

Fig. 1 shows a level scheme of the unbound  $^{15}\text{Ne}$ , the also unbound  $^{14}\text{F}$ , and the finally bound  $^{13}\text{O}$  [2]. In order to cast light on the decay mechanism of the ground state of  $^{15}\text{Ne}$  – be it a *diproton*, a *three-body*, or a *sequential* decay via a state in the unbound  $^{14}\text{F}$  – we studied its 3-body energy correlations and compared them to those in  $^{16}\text{Ne}$ , which is known to decay in a *three-body* way [3, 4, 5].




Fig. 2: Fractional relative energy ( $E_{pp}/E_{fpp}$ ) distributions in the ground states of  $^{15,16}\text{Ne}$  [2, 5]. See text for details.

Fig. 2 shows the ( $E_{pp}/E_{fpp}$ ) fractional relative energy

\* Work supported by EMMI, HIC for FAIR, the GSI-TU Darmstadt co-operation, BMBF, and NAVI. (B. J.) is a Helmholtz International Fellow.

of the ground states of  $^{15}\text{Ne}$  (filled diamonds), of  $^{16}\text{Ne}$  (open diamonds) and a *three-body* decay calculation for  $^{16}\text{Ne}$  [5] (full red line), and a calculation for *sequential* decay of  $^{15}\text{Ne}$  via the  $^{14}\text{F}$  ground state (black dashed line). The striking similarity to the pattern for  $^{16}\text{Ne}$ , combined with the discrepancy to the *sequential*-decay shape, leads us to conclude that, like in  $^{16}\text{Ne}$ , also the  $^{15}\text{Ne}$  ground state undergoes *three-body* decay.

Furthermore, we used the measured two-proton separation energy of  $^{15}\text{Ne}$  of 2.522(66) MeV, translated into an atomic mass excess of 40.215(69) MeV, to deduce the  $(1s_{1/2})^2$  occupation probability of its unbound valence-proton pair in the ground state. We followed the approach of Fortune [6] shown in Fig. 3, using a correlation between the  $(1s_{1/2})^2$  value for valence-nucleon pairs in  $Z = 8, 10$  mirror nuclei and their 2n-2p separation-energy difference in order to predict the  $^{15}\text{Ne}$  ground-state energy. Using our measured value of  $S_{2p} = 2.522(66)$  MeV, we have turned the relation around to predict an  $(1s_{1/2})^2$  content for the  $^{15}\text{Ne}$  ground state of 63(5) % (red square in Fig. 3).




Fig. 3: Relation of  $\Delta S_{2N}$  to  $P((1s_{1/2})^2)$  in  $Z = 8, 10$  mirror-nucleus pairs (based on [6]). See text for details.

## References

- [1] F. Wamers et al., GSI Report 2014-1, 136 p. (2014)
- [2] F. Wamers et al., Phys. Rev. Lett. **112**, 132502 (2014)
- [3] J. Marganec et al., Eur. Phys. J. A **51**: 9 (2015)
- [4] I. Mukha et al., Phys. Rev. C **77**, 061303 (R) (2008)
- [5] L.V. Grigorenko et al., Phys. Rev. Lett. **88**, 042502 (2002)
- [6] H.T. Fortune, Phys. Lett. B **718**, 1342 (2013)